



TCT@ACC-i2: Interventional Cardiology

PERIPHERAL ARTERY STENT THROMBOSIS: REPORT FROM THE EXCELLENCE IN PERIPHERAL ARTERY DISEASE (XLPAD) REGISTRY

Moderated Poster Contributions

TCT@ACC-i2 Moderated Poster Theater, Poster Hall B1

Sunday, March 15, 2015, 10:00 a.m.-10:10 a.m.

Session Title: TCT@ACC-i2: Interventional Cardiology III

Abstract Category: 31. TCT@ACC-i2: Carotid and Endovascular Intervention

Presentation Number: 2107M-05

Authors: Subhash Banerjee, Karan Sarode, Atif Mohammad, Osvaldo Gigliotti, Michael Luna, Tayo Addo, Mirza Baig, Shirling Tsai, Nicolas Shammass, Anand Prasad, Mazen Abu-Fadel, Andrew Klein, Emmanouil Brilakis, Veteran Affairs North Texas, Dallas, TX, USA

Background: There are limited data on infrainguinal peripheral artery stent thrombosis (ST).

Methods: We analyzed procedures performed between May 2005-March 2014 enrolled in the XLPAD registry. Occurrence of ST was determined by acute onset of lower limb pain and angiographic evidence of thrombus.

Results: ST occurred in 27 of 1,054 cases (2.6%) and was more likely in males (3.0% vs. 0.5%; $p=0.016$). Occurrence was similar in claudicants and in those with critical limb ischemia (2.2% vs. 2.8%; $p=0.823$). Compared to patients without ST, index procedures resulting in ST exhibited longer stent lengths (230.0 ± 103.1 mm vs. 109.3 ± 135.1 mm; $p<0.001$; Figure 1a), with more chronic total occlusions (88.9% vs. 52.2%; $p<0.001$; Figure 1b), fewer restenotic lesions (66.7% vs. 81.5%; $p=0.076$) and mostly superficial femoral artery (SFA) location (96.3% vs. 80.6%; $p=0.044$). ST was more frequent with drug-eluting stent (DES) implantation compared with bare-metal stents (7.5% vs. 2.5%; $p=0.060$). Dual antiplatelet therapy was equally prescribed (81.5% vs. 83.7%; $p=0.791$). Endovascular revascularization for ST tended to have higher procedural complication rates (8.3% vs. 4.5%; $p=0.299$).

Conclusion: Infrainguinal peripheral artery ST is more prevalent with DES implants in lesions with longer stent lengths and implanted for treatment of CTO predominantly in the SFA distribution.

